

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 June 2005 (02.06.2005)

PCT

(10) International Publication Number
WO 2005/049614 A2

- (51) International Patent Classification⁷: **C07D 475/04**, 487/14, A61K 31/505 // (C07D 487/14, 241:00, 239:00, 235:00)
- (74) Agent: **IZRAELEWICZ, Mark, H.**; Marshall, Gerstein & Borun LLP, 6300 Sears Tower, 233 South Wacker Drive, Chicago, IL 60606 (US).
- (21) International Application Number: PCT/US2004/038313
- (22) International Filing Date: 17 November 2004 (17.11.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/520,367 17 November 2003 (17.11.2003) US
60/520,368 17 November 2003 (17.11.2003) US
- (71) Applicants (for all designated States except US): **BIOMARIN PHARMACEUTICAL INC.** [US/US]; 105 Digital Drive, Novato, CA 94949 (US). **MERCK EPROVA AG** [CH/CH]; Im Laternenacker 5, CH-8200 Schaffhausen (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MOSER, Rudolf** [CH/CH]; Lahnhalde 11, CH-8200 Schaffhausen (CH). **GROEHN, Viola** [DE/CH]; Bahnhofstrasse 2, CH-8447 Dachsen (CH). **SCHUMACHER, Andreas** [DE/DE]; Am Neubrunnen 15, 79588 Efringen-Kirchen (DE). **MARTIN, Pierre** [CH/CH]; Meisenweg 38, CH-4310 Rheinfelden (CH).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published.
-- without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESSES FOR PREPARING TETRAHYDROBIOPTERIN, AND ANALOGS OF TETRAHYDROBIOPTERIN

(57) Abstract: Process for the preparation of tetrahydrobiopterin from neopterin and/or 6-substituted pterins with an improved yield and a high stereoselectivity. Also disclosed herein are novel individual intermediates prepared in the preparation of tetrahydrobiopterin, such as selectively protected neopterin useful for the preparation of tetrahydrobiopterin.

WO 2005/049614 A2

PROCESSES FOR PREPARING TETRAHYDROBIOPTERIN, AND ANALOGS OF TETRAHYDROBIOPTERIN

Cross-Reference To Related Application

This application claims the benefit under 35 U.S.C. § 119(e) of U.S.

- 5 Provisional Patent Applications Serial Nos. 60/520,367 and 60/520,368, both filed November 17, 2003, the disclosures of which are hereby incorporated herein by reference.

Background

Field of the Technology

- 10 The invention relates to a method for preparing tetrahydrobiopterin and related compounds and analogs of biopterin. More specifically, it relates to a stereoselective process for preparing 5,6,7,8-tetrahydro-6-(L-erythro-1',2'-dihydroxypropyl)pterin and includes the conversion of 6-(L-erythro-1,2,3-trihydroxypropyl)pterin and/or 6-substituted pterins to tetrahydrobiopterin.

15 Brief Description of Related Technology

- Tetrahydrobiopterin is a biogenic amine of the naturally-occurring pterin family. Pterins are present in physiological fluids and tissues in reduced and oxidized forms, however, only the 5,6,7,8-tetrahydrobiopterin is biologically active. Tetrahydrobiopterin is a chiral molecule, and the 6R enantiomer, and 1'R,2'S,6R
- 20 diastereomer of the tetrahydrobiopterin is the known biologically active form. The synthesis and disorders of tetrahydrobiopterin are described in Blau et al., Disorders of tetrahydrobiopterin and related biogenic amines, in Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Vogelstein B, eds. The Metabolic and Molecular Bases of Inherited Disease, 8th ed., New York: McGraw-Hill, 2001, at pages 1275-1776.

- 25 In a living body tetrahydrobiopterin plays a very important role as cofactor of essential enzymes (e.g., the aromatic amino acid hydroxylases, the nitric oxide synthetases, as a coenzyme in a catecholamine-serotonin synthesis.) Tetrahydrobiopterin is an indispensable compound for biosynthesis of the neurotransmitters dopamine and hydroxytryptamine, of noradrenalin, adrenaline, and

melatonin. The importance of tetrahydrobiopterin has been recognized in the course of the fundamental studies thereon. A deficiency of tetrahydrobiopterin causes serious neurological disorders like phenylketonuria (PKU) and Parkinson's disease.

Symptoms due to such diseases can be remarkably improved by administration of tetrahydrobiopterin. Further, it has been recognized that tetrahydrobiopterin is effective for curing infantile autism and depressions.

Such useful pharmacological activities, as well as the challenging chemical structures of the molecule, have stimulated many synthetic efforts directed toward the preparation of tetrahydrobiopterin. For example, tetrahydrobiopterin has been prepared by: (1) the reaction of 4-hydroxy-2,5,6-triaminopyrimidine (TAP) and 5-deoxy-L-arabinose as described in E. L. Patterson et al., J. Am. Chem. Soc., 78, 5868 (1956); (2) the reaction of TAP and 5-deoxy-L-arabinose phenylhydrazone, as described in Matsuura et al., Bull. Chem. Soc. Jpn., 48, 3767 (1975); (3) the reaction of TAP and triacetyloxy-5-deoxy-L-arabinose phenylhydrazone, as described in M. Viscontini et al., Helv. Chim. Acta., 60, 211 (1977); (4) the reaction of oxime and benzyl α -aminocyanoacetate and condensation of the resulting 3-(1,2-dihydroxypropyl)-pyrazine-1-oxide derivatives with guanidine followed by deoxygenation of the N-oxide, as described in E. C. Taylor et al., J. Am. Chem. Soc., 96, 6781 (1974); (5) the reaction of α -hydroxyketone (prepared from crotonic acid) and TAP, as described in M. Viscontini et al., Helv. Chim. Acta., 55, 574 (1972); and (6) the reaction of TAP having protected hydroxyl group and 4-acetoxy-2,3-epoxypentanal followed by oxidation with iodine and deprotection, as described in Matsuura et al., Chemistry of Organic Synthesis, Vol. 46, No. 6, p. 570 (1988), by protecting the hydroxyl group of S-alkyl lactate with a trityl group, reducing the resulting alkyl 2-trityloxypropionate to (S)-2-trityloxypropanol, oxidizing it to (S)-2-trityloxypropanal, treating it with a 2-furyl metal compound to form (1S, 2S)-1-(2-furyl)-2-trityloxy-1-propanal followed by oxidation and hydrolysis to form 2,3-dideoxy-6-trityloxyhepto-2-enopyranose-4-ulose, reducing it to 6-trityloxyhepto-2-ene-1,4,5-triol, acylating it to form 1,4,5-triacetyloxy-6-trityloxyhepto-2-ene followed by oxidation to afford 2,3 diacetyloxy-4-hydroxy-1-pentanal, treating it with phenylhydrazine to form a hydrazine, and condensing the hydrazine with a 3,5,6-

triaminopyrimidinol followed by oxidation and deacylation, as described in Japanese Kokai No. 221380/1989.

Each of these conventional processes for preparing tetrahydrobiopterin have several drawbacks, including, for example, expensive and sparsely available
5 carbohydrates are required as starting material to provide the asymmetric carbon atom at its side-chain, in that yield and purity are low due to multi-reaction steps, unstable intermediates are generated that require troublesome treatment operations, and troublesome purification procedures are required.

The prior processes for preparing tetrahydrobiopterin starting from 5-deoxy-L-
10 arabinose are economically disadvantageous, since 5-deoxy-L-arabinose of the required purity is only not readily available in large quantities. Also the product from the reactions involving 5-deoxy-L-arabinose is known to undergo degradation. Other prior preparation of tetrahydrobiopterin have the disadvantage that biopterin is produced in a DL-form and optical resolution is required for obtaining the desired L-
15 biopterin, thus leading to complicated process step and low yield. Indeed, in A. Kaiser, H.P. Wessel, *Helv. Chim. Acta*, Vol. 70, p. 766, 1987, states at page 768, that "These results and considerations demonstrate that no high-yield synthesis of biopterin from neopterin can be expected due to pyrrolo-pteridin formation upon activation of the side-chain terminus of neopterin."

20 Therefore, the conventional processes are unsuitable for industrial production of the compound and its derivatives. There exists a need for a process for the preparation of tetrahydrobiopterin, and analogs thereof in good yield using inexpensive starting material. A need also exists for an industrial scale process for the preparation of substantially optically pure tetrahydrobiopterin with an improved yield
25 and a high stereoselectivity.

Summary

One aspect of the processes and compounds described herein is a preparation of tetrahydrobiopterin from neopterin with an improved yield and a high stereoselectivity for the natural type (6R)-form. In one aspect, the method comprises

the steps of protecting the 2-amino group of neopterin with a 2-amino protective group, which may make the product more soluble, followed by carrying out a selective reaction on the primary hydroxyl group. In one embodiment, the primary hydroxyl group of neopterin is selectively protected with a primary hydroxyl protecting group, the secondary hydroxyl groups also selectively protected with a secondary hydroxyl protecting group, and reduction is carried out on the primary hydroxyl position in the side chain. In another embodiment, the primary hydroxyl group of neopterin is converted to its corresponding thioether, and reduction is carried out on this thioether. Deprotection of any of these groups may take place at any suitable time; for example, deprotecting the 2-amino group may occur immediately after selective protection of the primary hydroxyl group or may occur later. Alternatively, the method optionally includes erythro-selective reduction. The method optionally includes the step of hydrogenating and/or crystallizing tetrahydrobiopterin dihydrochloride.

Another aspect of the processes and compounds described herein is preparation of tetrahydrobiopterin from 6-substituted pterins including the steps of metalation of a protected 6-substituted pterin and coupling with lactic acid or a precursor of lactic acid. The method optionally includes the step of erythro-selective reduction of the 1'-keto group. Deprotection may take place at any suitable time; for example, immediately before erythro-selective reduction or afterwards. The method optionally includes the step of hydrogenating and/or crystallizing tetrahydrobiopterin dihydrochloride.

Yet another aspect of the present invention relates to novel individual intermediates, such as selectively protected pterin derivatives.

Further aspects and advantages of the invention will be apparent to those skilled in the art from a review of the following detailed description, taken in conjunction with the appended claims. While the processes and compounds are susceptible of embodiments in various forms, the description hereinafter includes specific embodiments of the invention with the understanding that the disclosure is illustrative, and is not intended to limit the inventions to the specific embodiments described herein.

Brief Description of the Drawings

Figure 1 is a schematic representation of the reaction scheme for preparing L-Neopterin.


Figure 2 is a schematic representation of a process described herein for the conversion of L-Neopterin to Tetrahydrobiopterin dihydrochloride salt.

Figure 3 is a schematic representation of a process described herein for the conversion of a 6-substituted Pterin to Tetrahydrobiopterin.

Figure 4 is a is a schematic representation of a process described herein for the conversion of L-Neopterin to Tetrahydrobiopterin, wherein the primary hydroxyl group on L-Neopterin is converted to its corresponding thioether, and the resulting thioether is then reduced to product a deoxygenated Neopterin derivative.


Detailed Description of the Preferred Embodiments

Tetrahydrobiopterin is a heterocyclic compound that performs a central role in a number of biological processes. The general structure of tetrahydrobiopterin is shown below:



Tetrahydrobiopterin contains three consecutive stereocenters, labeled above as 6, 1', and 2'. Tetrahydrobiopterin, like a number of biologically active molecules, exhibits a substantially heightened biological activity when a single stereoisomer and enantiomer. Described herein are processes for the preparation of a substantially single enantiomer and stereoisomer of tetrahydrobiopterin, and analogs thereof.

The substantially optically pure compound L-Neopterin (CAS No 2277-43-2) is used as the starting material in one embodiment of the processes described herein. The general structure of L-Neopterin is shown below:



A schematic representation of the process for preparing L-Neopterin from L-Arabinose is shown in Figure 1 (L-Neopterin is also available from Schircks Laboratories of Jona, Switzerland). The process for preparing L-Neopterin is also
 5 described in Pfeleiderer et al, *Helv. Chim. Acta*, Vol. 73, p. 808 (1990), and Viscontini et al, *Helv. Chim. Acta*, Vol. 53, p. 1202 1970, the disclosures of which are hereby incorporated herein by reference.

As used herein, the terms "linear chain alkyl" and "branch chain alkyl" encompasses, alkyl groups that may contain as few as one carbon atom or as many as
 10 fourteen carbon atoms, including but not limited to, cycloalkyl groups, methyl, ethyl, propyl, isopropyl, t-butyl, sec-butyl, cyclopentyl or cyclohexyl groups. The terms "linear chain alkyl" and "branch chain alkyl" also include alkyl groups that may be substituted with a variety of substituents, including but are not limited to, acyl, aryl, alkoxy, aryloxy, carboxy, hydroxy, carboxamido and/or N-acylamino moieties.

As used herein, an "aryl" encompasses, but is not limited to, a phenyl, pyridyl, pyrrol, indolyl, naphthyl, thiophenyl or furyl group, each of which may be substituted by various groups, which includes, but are not limited, acyl, aryl alkoxy, aryloxy, carboxy, hydroxy, carboxamido or N-acylamino moieties. Examples of aryloxy groups include, but are not limited to, a phenoxy, 2-methylphenoxy, 3-methylphenoxy and 2-
 20 naphthoxy. Examples of acyloxy groups include, but are not limited to, acetoxy, propanoyloxy, butyryloxy, pentanoyloxy and hexanoyloxy.

As used herein, the terms "alkoxycarbonyl", "acyl" and "alkoxy" encompass, but are not limited to, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, n-butoxycarbonyl, benzyloxycarbonyl, hydroxypropylcarbonyl, aminoethoxycarbonyl,
 25 secbutoxycarbonyl and cyclopentyloxycarbonyl. Examples of acyl groups include, but are not limited to, formyl, acetyl, propionyl, butyryl and penanoyl. Examples of

alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy, n-butoxy, sec-butoxy and cyclopentyloxy.

The solvent medium useful in the reactions of the processes described herein includes a wide variety of solvents. The reactions described herein are preferably performed wherein the reaction starting materials (*e.g.*, 6-substituted neopterin, neopterin, neopterin derivatives, biopterin and tetrahydrobiopterin) are dissolved in the solvent medium. Thus, the solvents used in the reactions described herein are preferably polar solvents capable of dissolving the polar compounds used and created according to the processes described herein. Preferably, the solvent is N,N-dimethylformamide (also referred to herein as DMF). Preferably, the concentration of the reactants in the reaction mixture is in the range of about 0.1% to about 20% by weight, more preferably 0.2% to 10%. Though the materials quickly dissolve in a polar reaction medium, at the beginning of a reaction the substances can exist in a solid form. In such a case, the substances can be gradually dissolved in the medium as the reaction proceeds.


One embodiment of the processes and compounds described herein includes a process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof from neopterin, including the following steps: (a) reacting the primary hydroxyl group of neopterin with a silyl protecting group; (b) protecting the secondary hydroxyl groups with a secondary hydroxyl protecting group; (c) converting the silyl group formed in step (b) to a surrogate group selected from the group consisting of halogens, sulfonates, and thioethers; (d) reduction at the substituted formed in step (e) to a methyl group; and (e) removing the secondary hydroxyl protecting group added at step (d). Step (c) can be performed by: i) direct conversion of the primary hydroxyl protecting group to a halogen; or (ii) selective cleavage of the silylether followed by a conversion of the protected primary hydroxyl group to a group selected from the group consisting of halogens, sulfonates, and thioethers. Preferably, the conversion in step (e) is performed by a direct conversion of the primary hydroxyl protecting group to a halogen. It has been found that this embodiment of the processes described herein can be performed without the protection of the 2-amino group on the neopterin. Under certain circumstances (*e.g.*, in order to avoid side reactions and degradation) it

may be preferable to first protect the 2-amino group of L-Neopterin before performing step (a) as described above. If the process of this embodiment is performed with the use of a 2-amino protecting group, the 2-amino protecting group is preferably removed after step (a) is performed. An example of the reactions of this process, wherein the 2-amino groups are protected/deprotected, is exemplified in Figure 2.

As exemplified in Figure 2 as Step 4, the 2-amino group can be protected before performing step (a). The protection of the 2-amino group on the L-Neopterin is preferably performing using a variety of protecting groups. Preferably, the protecting group for the 2-amino position on L-Neopterin is selected from the group consisting of dialkylformamidedialkylacetal groups, and pivaloyl groups. More preferably, the protecting group is one of N,N-dimethylformamidediethylacetal, and N,N-dimethylformamidedimethylacetal.

Protecting the 2-amino group dialkylformamidedialkylacetals yields the selective protection of the 2-amino group, and is described in Brederick et al, Chem. Ber., 101 41-50, (1968), and generally in Russel et al., Synlett 1992, p. 711, the disclosures of which are hereby incorporated herein by reference. Preferably, the reaction to protect the 2-amino group is carried out in a polar solvent, more preferably in dimethylformamide. In addition the 2-(N,N-dialkylaminomethylene-imino) Neopterin derivatives are much more soluble in non-polar organic solvents than the unprotected neopterin, and the protection of the 2-amino group to with a 2-(N,N-dialkylaminomethylene-imino) protecting group could be performed in a less polar solvent than DMF.

Another embodiment of the processes and compounds described herein is a compound of Formula 6, which is prepared according to Step 4:




6

wherein R1 is selected from the group consisting of single substituted linear chain

alkyl groups, single substituted branched chain alkyl groups, aryl substituted amido groups, an acetamido group, and a 2,2-dimethylpropanamido group. Preferably, R1 preferably single linear chain alkyl substituted alkylaminomethylene-imine groups, single branched chain alkyl substituted alkylaminomethylene-imine groups, double
 5 linear chain alkyl substituted alkylaminomethylene-imine groups, and double branched chain alkyl substituted alkylaminomethylene-imine groups.

Another protecting group that can be used to protect the 2-amino group is an acyl group, preferably a pivaloyl group. These compounds are obtained by the preparation of the acyl or tetrapivaloyl derivative of neopterin, followed by an alkaline
 10 hydrolysis of the three ester groups, as described in the literature, *e.g.*, Russell et al., Tet. Let., vol. 33, No. 23, pp 3371-3374 (1992), the disclosure of which is hereby incorporated herein by reference.


Another embodiment of the processes and compounds described herein is a 2-amino protected derivative of L-Neopterin of Formula 15:



wherein R5 is -COR', R' is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, aryl groups, and t-butyl; and R6 is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

20 The next step in the process, as exemplified in Figure 2 as Step 5, is the selective protection of the primary hydroxyl group of a compound of Formula 6 (as prepared according to Step 4), to yield a compound of Formulae 7 and 7a (both shown below).

25 Another embodiment of the processes and compounds described herein is a compound of Formula 7 (prepared according to Step 5):




7

- wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, and 2,2-dimethylpropanamide; and wherein R2 is a silyl group that is stable under acidic conditions. Preferably, R1 comprises N,N-dimethylaminomethylene amino, and R2 is selected from the group consisting of diethylisopropylsilyl, dimethylisopropylsilyl, dimethylphenylsilyl, diphenylisopropoxysilyl, diphenyl-t-butoxysilyl, di-t-butylmethylsilyl, di-t-butylsilylene, methyldiisopropylsilyl, methyldiphenylsilyl, t-butylmethoxyphenylsilyl, t-butyltrimethylsilyl, hexyldimethylsilyl, triethylsilyl, 1,1,3,3-tetra-isopropylidisiloxane, triisopropylsilyl, trimethylsilyl, trimethylsilyloxycabomyl, and t-butylphenylsilyl. More preferably, R2 is t-butylphenylsilyl.
- Other protecting groups and details of processes for their introduction/removal may be found by reference to Hart et al, J. Organic. Chem, 68(1) (2003), Corey et al, J. Am. Chem. Soc., 94(17) (1972), or Chaudary et al, Tet. Let., 2, pp 99-102 (1979), the disclosures of which are hereby incorporated herein by reference. The selective protection reaction is preferable carried out in a polar solvent, more preferably in dimethylformamide. The use of alkylchlorosilanes reagents as protecting agents yields in a highly selective protection of the primary hydroxyl function. According to this method there was no observed influence on the secondary hydroxyl groups within the molecule. The reaction is carried out in presence of a base, preferably the base is imidazole. The selective protection step can also be carried with a compound of Formula 15.

When a 2-amino protecting group is utilized in this embodiment, is possible to selectively cleave of the protection group at the 2-amino function with an *in situ*


reaction. When the 2-amino group is protected as an acyl group, the selective deprotection can be performed with ammonium hydroxide in dioxane, when the 2-amino group is protected as a linear or branched chain alkyl substituted sulfur group, the selective deprotection can be performed with ammonium hydroxide, when the 2-amino group is protected as an N,N-dimethylaminomethylene substituted amino group, the selective deprotection can be performed with ammonium hydroxide in dioxane and preferably with zinc dichloride in ethanol. The selective deprotection of the 2-amino group yields Formula 7a.

Thus, another embodiment of the processes and compounds described herein is a compound of Formula 7a:



wherein R₂ is a silyl group that is stable under acidic conditions. Preferably, R₂ is selected from the group consisting of diethylisopropylsilyl, dimethylisopropylsilyl, dimethylphenylsilyl, diphenylisopropoxysilyl, diphenyl-t-butoxysilyl, di-t-butylmethylsilyl, di-t-butylsilylene, methyl-diisopropylsilyl, methyl-diphenylsilyl, t-butylmethoxyphenylsilyl, t-butyl-dimethylsilyl, t-hexyldimethylsilyl, triethylsilyl, 1,1,3,3-tetra-isopropyl-disiloxane, triisopropylsilyl, trimethylsilyl, trimethylsilyloxycabomyl, and t-butyl-diphenylsilanoyl. More preferably, R₂ is t-butyl-diphenylsilane

The next step in the process, as exemplified in Figure 2 as Step 6, is the protection of the secondary hydroxyl groups of a compounds of Formulae 7 and/or 7a (as prepared according to example shown in Step 5), to yield a compound of Formula 8 (shown below).



8

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R2 is a silyl group that is stable under acidic conditions; and R4 a substituted acetal or ketal group that is stable under alkaline conditions. Preferably, R4 is a substituted acetal or ketal group is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals. More preferably, R4 is selected from the group consisting of methylene acetal, ethylidene acetal, t-butylmethylidene ketal, 1-t-butylethylidene ketal, 1-phenylethylidene ketal, 1-(4-methoxyphenyl)ethylidene acetal, 2,2,2-trichloroethylidene acetal, acrolein acetal, cyclopentylidene ketal, cyclohexylidene ketal, cycloheptylidene ketal, benzylidene acetal, p-methoxybenzylidene acetal, 2,4-dimethoxybenzylidene ketal, 3,4-dimethoxybenzylidene acetal, 2-nitrobenzylidene acetal, 4-nitrobenzylidene acetal, mesitylene acetal, 1-naphthaldehyde acetal, benzophenone ketal, and isopropylideneketal.


Cyclic ortho-esters and other 1,2-diol protective groups which are stable to alkaline conditions and cleaved under acidic conditions are suitable protecting groups for the secondary hydroxyl groups.

The reaction of exemplified in Figure 2 as Step 6 is preferably performed a polar solvent, more preferably in acetone. Also preferable, R4 is acetonedimethylacetal, and the reaction is performed in acetone and in the presence of p-toluenesulfonic acid. Other protecting groups and details of processes for their introduction/removal may be found in "Protective Groups in Organic Synthesis",

Green et al., 3rd Ed. (1999) Wiley & Sons, p 201-245, the disclosure of which is hereby incorporated herein by reference.

The next step or series of steps in the process, as exemplified in Figure 2 as Step 7, and Step 9 and Step 7a conversion of the silyl ether to a halide. As shown, for example in Figure 2, Step 7, the silyl ether (such as in a compound of Formula 8) is directly converted to a halide, and exemplified in Steps 9 and 7a of Figure 2, the conversion goes through a deprotection step. Through either pathway, the result is the formation of a compound of Formula 9 (shown below).

Thus, another embodiment of the processes and compounds described herein is a compound of Formula 9:




9

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and wherein R5 is a halogen.

The halogen is preferably introduced into the molecule by triphenylphosphine halogen, preferably with triphenylphosphine bromide. The reaction is preferably carried out a solvent selected from the group consisting of dichloromethane, dimethylformamide, and dimethylacetamide. Other reaction conditions and details of the conversion of the silyl ether to a halide can be found in Hanessian et al, J. Org. Chem., 34(7), p 2163 (1969), Kim et al, J. Org. Chem., 53, p 3111-3113 (1988), Ashton, J. Org. Chem., 61(3), p905 (1996), Aizpurua et al, J. Org. Chem., 51(25), p

4942 (1986), and Mattes, Tet. Let., 28(15), p 169 (1987), the disclosures of which are hereby incorporated herein by reference.

As described above, an alternative method for the preparation of a compound of Formula 9 includes the deprotection of the primary hydroxyl followed by the formation of the halide. As exemplified in Figure 2 as Step 9, when a compound of Formula 8 is treated with a base in a protic solvent (e.g., KOH in CH₃OH), the silyl ether protecting group is cleaved to form a compound of Formula 11. Thus, another embodiment of the processes and compounds described herein is a compound of Formula 11:




wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; and R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals.

The deprotection of the primary hydroxyl group as described above (e.g., as exemplified in Step 9 of Figure 2) is preferably carried out in an alcohol, more preferably in methanol. The conversion may also be achieved by using fluorides (e.g., tetrabutylammoniumfluoride in tetrahydrofuran or other apolar solvents).

The deprotected primary hydroxyl group is then converted to a halide as shown, for example, in Step 7a in Figure 2. This conversion is performed under the same conditions as described above for example shown in Step 7 of Figure 2, and yields a compound of Formula 9. Further details regarding the reaction conditions and

processes for this conversion can be found in "Comprehensive Organic Transformations", R.C. Larock, 2nd ed., Wiley VCH, p689-697 (1999), the disclosure of which is hereby incorporated herein by reference.


As an alternative to the preparation of a compound of Formula 9, the primary hydroxyl group on a compound of Formula 11 can be converted to a sulfonate group (e.g., a tosylate group) such as in a compound of Formula 12, such as, for example, as shown in Step 10 of Figure 2. Thus, another embodiment of the processes and compounds described herein is a compound of Formula 12:



12

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and R6 is selected from the group consisting of linear chain alkyl substituted sulfonates, branched chain alkyl substituted sulfonates, and aryl substituted sulfonates. Preferably, R4 is dimethylacetal, R3 is an N,N-dimethylaminomethylene substituted amino group, and R6 is a tosyl group. The reaction exemplified in Step 10 of Figure 2 is preferably carried out with sulfonylchloride in the presence of a base.

The next step in the process is the preparation of a compound of Formula 10. Both compounds of Formulae 9 and 12 can be converted to a compound of Formula 10. As an example, these conversions are exemplified in Figure 2 as Step 11, and Step 8. Thus, another embodiment of the processes and compounds described herein is a compound of Formula 10:




10

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; and R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals. Preferably, R4 is dimethylacetal, and R3 is an N,N-dimethylaminomethylene substituted amino group.

The reduction of a compound of Formulae 12 and 9 is preferably carried out with sodium borohydride and/or lithium aluminum hydride, and preferably in a polar, non-aqueous aprotic solvent such as dimethyl sulfoxide, dimethylformamide or sulfolane. Other reaction conditions and details of regarding this step can be found in "March's Advanced Organic Chemistry, Reaction Mechanisms and Structures" by Smith and March, 5th edition, Wiley & Sons, Inc., p. 524-526 (2001), European Patent No. 0 165 595 A2, Zinner et al, Chem. Ber., 92,1618 (1959), Zinner et al, Chem. Ber., 92, 2893 (1959), Bell et al, J. Org. Chem., 34, 3923 (1969), Hutchins et al, Tel. Let., 3495 (1969), and Hutchins et al, J. Org. Chem., 24, 2923, (1969), the disclosures of which are hereby incorporated herein by reference.

As an alternative of the preparation of process a halide of Formula 9 and a tosylate of Formula 12, a sulfonate can be prepared from the deprotected alcohol of Formula 11. As described below, the sulfonate can be converted into its corresponding thioether with the use of the Mitsunobu reaction, and then the resulting thioether can be reduced to yield a compound of Formula 10. Thus, another embodiment of the processes and compounds described herein is a compound of Formula 11a:




11a

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and R7 is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

This reaction to convert a compound of Formula 11a into a compound of Formula 10 is preferably performed using the Mitsunobu reaction (*e.g.*, PPh₃/ROOCN=NCOOR), followed by reduction of the resulting thioether using, for example, Raney-Nickel. Preferably, the reduction of the thioether is performed using a Raney-Nickel and hydrogen, more preferably Raney-Nickel, hydrogen, in an ethanol solvent medium.

The next step in the process, as exemplified in Figure 2 as Step 12, is the reduction of a compound of Formula 10 (as prepared according to Steps 8 and 12), to stereoselectively yield tetrahydrobiopterin. The tetrahydrobiopterin can then be converted to its salt form, including but not limited to its dihydrochloride salt as shown below:



The reduction of a compound of Formula 10 is carried out either according to

methods well known in the common literature (*e.g.*, sodium borohydride in a alkaline medium) or preferably with a catalytic amount of platinum dioxide and hydrogen.

Tetrahydrobiopterin may be isolated preferably as dihydrochloride by crystallization techniques well known in the art, such as suspension, precipitation, re-crystallization, evaporation, solvent like water sorption methods or decomposition of solvates. Diluted, saturated, or super-saturated solutions may be used for crystallization, with or without seeding with suitable nucleating agents.

Another embodiment of the processes and compounds described herein is a process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof, including the following steps: (a) reacting pterin at the C-6 position to prepare a 6-substituted pterin; (b) protecting the primary amine group at C-2 of neopterin with a 2-amino protecting group; (c) metalation of the protected 6-substituted pterin; (d) coupling of the product of the metalation of the protected 6-substituted pterin with lactic acid or a precursor of lactic acid; (e) removing the 2-amino protecting group; and (f) erythro-selective reduction.


The first step in this embodiment utilizes a 6-substituted pterin, including but not limited to 6-halogenated pterins and 6-sulfonated pterins. Preferably, the 6-substituted pterins are 6-halogenated pterins, more preferably, the pterin starting material is selected from the group consisting of 6-chloropterin, 6-bromopterin, and 6-iodopterin. It has been found that 6-iodopterin is the preferred 6-halogenated pterins for use in the coupling reaction described below.

The first step in this embodiment is the protection the 2-amino group of the 6-substituted pterin, as exemplified in Figure 3 as Step 1, is the protection of the 2-amino group in the 6-substituted pterin. The protection of the 2-amino group is performed as described above (for the preparation of a compound of Formula 6 (exemplified as Step 4 in Figure 2), and the product of this step is a compound of Formula 2 (shown below).

Preferably, the protecting group used to protect the 2-amino group is selected from the group consisting of linear chain alkyl single substituted amido groups,

branched chain alkyl single substituted amido groups, aryl substituted amido group, a pivaloyl group, and 2,2-dimethylpropanamido. More preferably, the protecting group is a pivaloyl group.

Thus, another embodiment of the processes and compounds described herein
5 is a compound of Formula 2:




2

wherein X is selected from the group consisting of chlorine, bromine, iodine, and sulfonates; R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups,
10 double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, single linear chain alkyl substituted alkylaminomethylene-imine groups, single branched chain alkyl substituted alkylaminomethylene-imine groups, double linear chain alkyl
15 substituted alkylaminomethylene-imine groups, and double branched chain alkyl substituted alkylaminomethylene-imine groups; and R2 is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

The next step in this embodiment of the processes disclosed herein is the
20 metalation of the 6-substituted pterin as exemplified in Figure 3 as Step 2. Preferably, the metalation of the protected 6-substituted pterin is performed with a reagent selected from the group consisting of RMgX (*i.e.*, a Grignard reagent), alkyl-metal complexes, and metals, wherein X is a halogen, and R is selected from the group consisting of alkyl groups, and aryl groups. Preferably, the alkyl-metal complex is an
25 alkyl-metallic lithium complex, more preferably n-butyllithium and/or t-butyllithium.

The metalation at the 6-position on the pterin produces a pterin derivative, which is not isolated according to this embodiment, but exists for a substantial period of time. Thus, another embodiment of the processes and compounds described herein is a compound of Formula 3:



wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups; branched chain alkyl substituted sulfur groups, and 2,2-dimethylpropanamide; R2 is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups; and M is selected from the group consisting of boron, silicon, zirconium, titanium, sodium, aluminum, nickel, cobalt, scandium, chromium, ytterbium, lithium, magnesium, zinc, palladium, copper, manganese, cesium, and tin.

10

15

The metalation reaction is preferably performed in non-polar solvents such as ethers, preferably diethylether, dioxane, and/or tetrahydrofuran (THF). Preferably, the metalation is performed with a Grignard reagent, and preferably the Grignard reagent is isopropylmagnesiumchloride. The reaction temperature during the metalation set is preferably kept in the range of about -80°C up to about +30°C, and preferably one to four equivalents of the metalating reagent (*e.g.*, Grignard reagent) are used for the metalation.


20

The next step in this embodiment is the coupling of the product from the metalation step with lactic acid or a lactic acid precursor as exemplified in Figure 3 as Step 3. Preferably, the coupling is performed between the protected 6-metalated pterin and a protected lactic acid chloride, more preferably between the protected 6-

25

metalated pterin and a hydroxyl protected lactic acid chloride such as 2-acetoxypionic chloride. Preferably, when a precursor of lactic acid is used in this step, the precursor of lactic acid is selected from the group consisting of 2-oxopropanoyl chlorides, and 2-oxopropanal.

- 5 It has been found to be preferable that the metalation step and the coupling step can be performed in the same reaction vessel. Accordingly, another embodiment of the processes and compounds described herein is a compound of Formula 3:




4

- 10 wherein R₁ is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R₂ is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups; and R₃ is an acyl group.
- 15 Preferably, R₁ is an N,N-dimethylaminomethylene substituted amino group.


- The coupling step can also be performed with a 2-oxopropanoyl chlorides or with 2-oxopropanale with the use of Pd(OAc)₂, Me₆Sn₂, PPh₃ in dioxane, then Pd(PPh₃)Cl₂ and lactic acid chloride or derivatives thereof. A discussion of the reaction conditions and details of processes related to the coupling reaction can be found in Bradshaw et al, J. Chem. Soc., Perkin Trans., 1, 3239-3244 (2001), and Pellicciari et al, Tetrahedron Lett., 3003-3004 (1992), the disclosures of which are hereby incorporated herein by reference. It has also been found preferable that diketones are formed when these reactants are used in the coupling step. Thus,
- 25 preferably when diketones are formed, the coupling step is followed by the further step of reducing the resulting diketones.

The next step in this embodiment of the processes described herein is deprotecting of the coupling product, and the erythro-selective reduction of the deprotected product to yield tetrahydrobiopterin as exemplified in Figure 3 as Steps 4 and 5. It has been found that when an acyl group is used as the protecting group R3 in a compound of Formula 4, the protecting group on the can be performed wherein on the protecting group at the 2-amino position is removed. Thus, another embodiment of the processes and compound described herein is a compound of Formula 5:



wherein R3 is an acyl group.

The tetrahydrobiopterin prepared by the reduction, as exemplified in Figure 3 as Step 5, can then be converted to its salt form, including but not limited to its dihydrochloride salt as shown below:



The erythro-selective reduction of a compound of Formula 4 is carried out either according to methods well known in the common literature (*e.g.*, sodium borohydride in a alkaline medium) or preferably with a catalytic amount of platinum dioxide and hydrogen.

Tetrahydrobiopterin may be isolated preferably as dihydrochloride by crystallization techniques well known in the art, such as suspension, precipitation, recrystallization, evaporation, solvent like water sorption methods or decomposition of solvates. Diluted, saturated, or super-saturated solutions may be used for crystallization, with or without seeding with suitable nucleating agents.

It has been found that the primary hydroxyl of neopterin can be reduced, and leave a methyl group at the C-3' position of the neopterin. This conversion is performed by converting the primary hydroxyl to a thioether, and then reduction of the thioether to a methyl group. Thus, another embodiment of the processes and compounds described herein is a process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof from neopterin, including the following steps: (a) protecting the primary amine group at C-2 of neopterin with a 2-amino protecting group; (b) converting the primary hydroxyl group of neopterin to a thioether; and (c) reduction of the thioether leaving a methyl group at the C-3' position on the neopterin side chain.

Quite surprisingly, it has been found that when step (c) above is performed at elevated temperatures (*e.g.*, above 50 degrees Celsius), the reaction conditions for the reduction reaction of step (c) will also remove the 2-amino protecting group and perform an erythro-selective reduction of the C5-C6 and C7-C8 double bonds on neopterin to yield tetrahydrobiopterin. Thus, it is preferred to perform the reduction reaction of step (c) at elevated temperatures and allow the reaction to be performed under an excess of reducing agent to allow the removal of the 2-amino protecting group and an erythro-selective reduction.

As described below, if the reduction of step (c) does not result in the erythro-selective reduction of the C5-C6 and C7-C8 and the removal of the 2-amino protecting group, the process further includes the two additional steps of removal of the 2-amino protecting group and an erythro-selective hydrogenation.

It has been found that the purity and stability of the tetrahydrobiopterin product produced according to this embodiment can be improved by forming a salt of tetrahydrobiopterin. Thus, the tetrahydrobiopterin is preferably conferred to its corresponding salt, more preferable to the dichloride salt of tetrahydrobiopterin. Preferably, the dichloride salt of tetrahydrobiopterin is further recrystallized.


The first step in this embodiment, as exemplified in Figure 4 as Step 1, is the protection of the 2-amino group on L-Neopterin. The protection of the 2-amino group on the L-Neopterin is preferably performing using a variety of protecting groups.

Preferably, the protecting group for the 2-amino position on L-Neopterin is selected from the group consisting of dialkylformamidedialkylacetal groups, and pivaloyl groups. More preferably, the protecting group is one of N,N-dimethylformamidediethylacetal, and N,N-dimethylformamidedimethylacetal.

5 Protecting the 2-amino group dialkylformamidedialkylacetals yields the selective protection of the 2-amino group, and is described in Brederick et al, Chem. Ber., 101 41-50, (1968), and generally in Russel et al., Synlett 1992, p. 711, the disclosures of which are hereby incorporated herein by reference. Preferably, the reaction to protect the 2-amino group is carried out in a polar solvent, more preferably
10 in dimethylformamide. In addition the 2-(N,N-dialkylaminomethylene-imino) Neopterin derivatives are much more soluble in non-polar organic solvents than the unprotected neopterin, and the protection of the 2-amino group to with a 2-(N,N-dialkylaminomethylene-imino) protecting group could be performed in a less polar solvent than DMF.

15 The second step in this embodiment, as exemplified in Figure 2 as Step 2, is the conversion of the primary hydroxyl group to a thioether. This conversion is preferably performed with the use of the Hata reagent. Thus, the primary hydroxyl is selectively converted to a thioether with the use of a disulfide reagent and a trialkylphosphine reagent, more preferably diphenyl disulfide and tributylphosphine.

20 Another embodiment of the processes and compounds disclosed herein is a compound of Formula 15:



15

wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups,
25 double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, single linear

chain alkyl substituted alkylaminomethyleneimine groups, or single branched chain alkyl substituted alkylaminomethyleneimine groups, double linear chain alkyl substituted alkylaminomethyleneimine groups, double branched chain alkyl substituted alkylaminomethyleneimine groups; and R2 is selected from the group
5 consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups. Preferably, R1 is a dialkylalkylaminomethyleneimine group, more preferably, dimethylaminomethyleneimine. Also preferably, R2 is benzene.

The next step in the process of this embodiment, as exemplified in Figure 4 as Step 3, is the reduction of the thioether, wherein the net result is the replacement of
10 the thioether with a hydrogen (*i.e.*, 2-amino protected Bioplerin). Preferably, the reduction of the thioether is performed with the use of a Raney-Nickel a reducing agent. It has been found that the reduction of the thioether does not proceed to yield product (*i.e.*, 2-amino protected L-Bioplerin) when the reaction is carried out in a protic solvent. Thus, the reduction reaction is preferably performed in a polar aprotic
15 solvent and at room temperature.

As described above, when the reduction of the thioether is performed at room temperature and carefully monitored, the reduction does not result in a removal of the 2-amino protecting group and an erythro-selective reduction. Accordingly, the next two steps in this embodiment, as exemplified in Figure 4 as Steps 4 and 5, include the
20 deprotection of the 2-amino group (*i.e.*, removal of the 2-amino protecting group), and the erythro-selective reduction of the product resulting from the deprotection (*i.e.*, L-Bioplerin). The specific reaction conditions for the deprotection and reduction steps are described above, and in the references listed above.

EXAMPLES

25 The following examples are provided to illustrate the processes and compounds described herein but are not intended to limit the scope of the processes and compounds described herein.

Example 1

The preparation of a N2-N,N-Dimethylaminomethylene-L-neopterin (a

compound of Formula 6 wherein R1 is an dimethylaminomethylene-imine group) was prepared by adding 15.8 ml of N,N-dimethylformamidediethylacetal to a flask containing a suspension of 11.68 g of L-neopterin and 850 ml of dry N,N-dimethylformamide. The mixture was stirred at room temperature until all starting material dissolved. After 6 hours of stirring at room temperature, 280 ml dry methanol was added and the reaction mixture and the mixture was stirred for an additional 12 hours. After evaporating all solvents the residue from the reaction mixture was suspended in 500 ml of dichloromethane, the suspended material was then collected and washed with 150 ml of dichloromethane, and the resulting product was vacuum dried at 40°C to give 11.23 g of N²-N,N-dimethylaminomethylene-L-neopterin.

The ¹H-NMR-data (200 MHz, solvent: DMSO-d₆) for the protected L-Neopterin is as follows: 11.98 ppm, bs, N³-H; 1H, 8.79 ppm, s, CH=N, C⁷-H, 2H; 5.64 ppm, d, C^{1'}-OH, 1H; 4.75 ppm, d, C^{2'}-OH, 1H; 4.63 ppm, dd, C^{1'}-H, 1H; 4.47 ppm, t, C^{3'}-OH, 1H; 3.81 ppm, m, C^{2'}H, 1H, 3.54 ppm, m, C^{3'}H₁, 1H; 3.43 ppm, m, C^{3'}H₂, 1H; 3.22 ppm, s, N-CH₃, 3H; 3.09 ppm, s, N-CH₃, 3H.

Example 2

The selective protection of the primary hydroxyl group was performed with the 2-amino protected L-Neopterin that was prepared according to Example 1. The N²-N,N-Dimethylaminomethylene-3'O(t-butyl-diphenylsilyl)-L-neopterin was prepared by first suspending 10 g of N²-N,N-Dimethylaminomethylen-L-neopterin in 250 ml of dry N,N-dimethylformamide, and then adding 4.9 g imidazole and 10g t-butyl-diphenylchlorosilane to the reaction mixture. After the reaction mixture stirred at room temperature for 2 hours an additional 0.5 g imidazole and 1 g t-butyl-diphenylchlorosilan. After the reaction mixture stirred for an additional 14 hours at room temperature the reaction mixture was evaporated to dryness and the residue was purified by flash chromatography on silica gel using a gradient of dichloromethane/methanol of 9:1 to 6:4. Product fractions were pooled and evaporated. The residue was suspended in 100 ml of isopropanol and the product was collected, washed with isopropanol, and vacuum dried at 40°C to give 9.3 g of N²-N,N-Dimethylaminomethylen-3'(t-butyl-diphenylsilyl)-L-neopterin.

The ¹H-NMR-data (200 MHz, solvent: DMSO-d₆) for the product is as follows: 11.99 ppm, bs, N3-H; 1H; 8.82 ppm, s, CH=N, C7-H, 2H; 7.63 ppm, m, Ph, 4H; 7.41 ppm, m, Ph, 6H; 5.73 ppm, d, C1'-OH, 1H; 5.00 ppm, d, C2'-OH, 1H; 4.79 ppm, dd, C1'-H, 1H; 4.05 ppm, m, C2'H, 1H, 3.78 ppm, m, C3'H1, 1H; 3.68 ppm, m, C3'H2, 1H; 3.23 ppm, s, N-CH₃, 3H; 3.10 ppm, s, N-CH₃, 3H; 0.93 ppm, s, C(CH₃)₃, 9H.

Example 3

The selective protection of the primary hydroxyl group was also performed with the 2-amino protected L-Neopterin that was prepared according to Example 1, and after the selective protection, the deprotection of the 2-amino group was performed in the same reaction flask to yield 3'O-(t-butyl-diphenylsilyl)-L-neopterin.

To a suspension of 8 g of N₂-N,N-Dimethylaminomethylen-L-neopterin in 200 ml of dry N,N-dimethylformamide were added 3.92 g imidazole and 8g t-butyl-diphenylchlorosilane. After the reaction mixture stirred for 2 hours at room temperature, an additional 0.4 g of imidazole and 0.8 g of t-butyl-diphenylchlorosilane was added. The reaction mixture was then allowed to stir at room temperature for 14 hours after which the reaction mixture was evaporated to dryness and the residue of crude N₂-N,N-Dimethylaminomethylen-3'(t-butyl-diphenylsilyl)-L-neopterin was dissolved in 160 ml ethanol. Upon dissolution in ethanol, 15 g of zinc-chloride was added to the reaction flask, and the mixture was heated to 80°C for 3 hours. During the course of the three hours a solid separated out from the mixture. The suspension was then cooled to 58°C and the solid was collected, washed with 100 ml ethanol and dried in vacuum at 40°C to give 5 g of 3'(t-butyl-diphenylsilyl)-L-neopterin. Another 5g fraction of 3'(t-butyl-diphenylsilyl)-L-neopterin was obtained from the filtrate after leaving it at room temperature for 24 hours.

The ¹H-NMR-data (200 MHz, solvent: DMSO-d₆) for 3'O-(t-butyl-diphenylsilyl)-L-neopterin was as follows: 11.40 ppm, bs, N3-H; 1H, 8.73 ppm, s, CH=N, C7-H, 2H; 7.63 ppm, m, Ph, 4H; 7.42 ppm, m, Ph, 6H; 6.86 ppm, bs, NH₂, 2H; 5.68 ppm, d, C1'-OH, 1H; 4.97 ppm, d, C2'-OH, 1H; 4.74 ppm, dd, C1'-H, 1H; 4.02 ppm, m, C2'H, 1H, 3.77 ppm, m, C3'H1, 1H; 3.66 ppm, m, C3'H2, 1H; 0.93 ppm,

s, C(CH₃)₃, 9H.

Example 4

The protection of the secondary hydroxyl groups was performed to prepare 1'2'-isopropylidene-3'O-(t-butyl-diphenylsilyl)-L-neopterin by adding 3.8 g para-
5 toluenesulfonic acid to a reaction flask containing 10g 3'(t-butyl-diphenylsilyl)-L-neopterin (prepared according to Example 3) in 50 ml acetone-dimethylacetal. The reaction mixture was allowed to stir for 14 hours at room temperature. The resulting solid was collected, washed with 30 ml of acetone-dimethylacetal and vacuum dried at 35°C to yield 6.5 g of 1'2'-isopropylidene-3'(t-butyl-diphenylsilyl)-L-neopterin.

10 The foregoing description is given for clearness of understanding only, and no unnecessary limitations should be understood therefrom, as modifications within the scope of the processes and compounds described herein may be apparent to those having ordinary skill in the art.

Example 5

15 The conversion of the primary hydroxyl of L-Neopterin to a thioether was performed to prepare 6-((1R,2R)-1,2-dihydroxy-3-phenylthiopropyl)-2-[(methylethyl)amino]-3-hydropteridin-4-one (the compound labeled "15" in Figure 4). To a flask containing 50 grams of 6-((2S,1R)-1,2,3-trihydroxypropyl)-2-[(1Z)-1-aza-2-(dimethylamino)vinyl]-3-hydropteridin-4-one (DMA-Neopterin, the compound
20 labeled "6" in Figure 2) and 400 ml of dimethylaminoacetate was added to the flask. To this stirring mixture 1 molar equivalent of diphenyl disulfide and tributylphosphine were added to the flask. It was discovered that the DMA-Neopterin has a low solubility in dimethylaminoacetate at room temperature (approximately 3.5 mg/ml). It is believed that the thioether product is more soluble in dimethylaminoacetate. The
25 reaction mixture was allowed to was allowed to stir for 4 hours at room temperature. Nine grams of the thioether product was isolated.

Example 6

The thioether product prepared in Example 5 was reduced with Raney-Nickel according to the following procedure. The thioether (9 grams, 22 mmole) was added

to a flask, and the flask was charged with 360 ml of ethanol. To a stirring mixture of the thioether in ethanol, 90 grams of Raney-Nickel in ethanol, and the reaction mixture was placed under an atmosphere of hydrogen of 5 bar of pressure. The reaction mixture was allowed to stir for 17 hours.

- 5 The hydrogen pressure was then released, the Raney-Nickel was filtered off from the reaction mixture, and 2.25 ml of concentrated hydrochloric acid was added to the filtrate. The resulting dihydrochloride salts were then collected and separated. Two diastereomers of BH₄ dihydrochloride salt was formed in the reaction, the desired (6R) form was prepared in 15.8 percent yield, and the undesired (6S) form
- 10 was prepared in 9.3% yield.

The foregoing description is given for clearness of understanding only, and no unnecessary limitations should be understood therefrom, as modifications within the scope of the processes and compounds described herein may be apparent to those having ordinary skill in the art.

WHAT IS CLAIMED IS:

1. A process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof from neopterin, comprising the following steps:
 - (a) reacting the primary hydroxyl group of neopterin with a silyl protecting
5 group;
 - (b) protecting the secondary hydroxyl groups with a secondary hydroxyl protecting group;
 - (c) converting the silyl group formed in step (b) to a surrogate group selected from the group consisting of halogens, sulfonates, and thioethers;
 - 10 (d) reduction at the substituted formed in step (e) to a methyl group; and
 - (e) removing the secondary hydroxyl protecting group added at step (d).
2. The process of claim 1, further comprising the step of protecting the primary amine group at C-2 of neopterin with a 2-amino protecting group before performing said step (a).
- 15 3. The process of claim 2, further comprising the step of removing the 2-amino protecting group after step (a) is performed.
4. The process of claim 1, further comprising the step of isolating the tetrahydrobiopterin product after step (e) is performed.
5. The process of claim 1, further comprising the step of erythro-selective
20 hydrogenation of the product resulting from step (e).
6. The process of claim 5, wherein said erythro-selective hydrogenation is performed by a reducing agent selected from the group consisting of sodium

borohydride, and platinum dioxide and hydrogen.

7. The process of claim 2, wherein said 2-amino protecting group said 2-amino protecting group comprises a protecting group selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, single substituted amino groups, double substituted amino groups, aryl single substituted amino groups, linear alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, linear alkyl single substituted alkylaminomethylene-imine groups, linear alkyl double substituted alkylaminomethylene-imine groups, branched alkyl single substituted alkylaminomethylene-imine groups, and branched alkyl double substituted alkylaminomethylene-imine groups.

8. The process of claim 7, wherein said 2-amino protecting group comprises a dialkylformamidedialkylacetal group.

9. The process of claim 8, wherein said dialkylformamidedialkylacetal group is selected from the group consisting of N,N-dimethylformamidediethylacetal, and N,N-dimethylformamidedimethylacetal.

10. The process of claim 7, wherein said 2-amino protecting group comprises a pivaloyl derivative of neopterin.


11. The process of claim 10, wherein said protecting the 2-amino group of neopterin is prepared by a two step process comprising, preparing a tetrapivaloyl derivative of neopterin, and performing alkaline hydrolysis on three pivaloyl groups on said tetrapivaloyl derivative of neopterin.

12. The process of claim 1, wherein said step (c) is performed by a direct conversion of the primary hydroxyl protecting group to a halogen.

13. The process of claim 1, wherein said step (c) is performed by selective

cleavage of said silylether formed in step (a) followed by a conversion of the resulting primary hydroxyl group to a group selected from the group consisting of halogens, sulfonates, and thioethers.

14. The process of claim 2, wherein the product resulting from said step (a) comprises a compound of the Formula 20:



- wherein R5 is -COR', R' is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, aryl groups, and t-butyl; and R6 is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

15. The process of claim 1, wherein said silyl protecting group comprises a protecting group selected from the group consisting of linear chain alkyl substituted silyl groups, branched chain alkyl substituted silyl groups, and aryl substituted silyl groups.
16. The process of claim 1, wherein said silyl protecting group comprises a silyl protecting group that is stable under acidic conditions.

17. The process of claim 15, wherein said silyl protecting group is selected from the group consisting of t-butyldimethylsilane, and t-butyldiphenylsilane.

18. The process of claim 1, wherein said secondary hydroxyl protecting group comprises an acetal or a ketal.

19. The process of claim 18, wherein said acetal or said ketal comprises isopropylideneketal.
20. The process of claim 1, wherein said silyl protecting group is exchanged with a halogen.
- 5 21. The process of claim 20, wherein said halogen comprises bromine.
22. The process of claim 20, wherein said exchange of said silyl protecting group is performing using a triphenylphosphine halogen.
23. The process of claim 1, wherein said silyl protecting group is selectively cleaved off by alkaline hydrolysis and the product of said hydrolysis is
10 transformed by halogenation.
24. The process of claim 23, wherein said halogenation is performed with a triphenylphosphine halogen.
25. The process of claim 24, wherein said triphenylphosphine halogen comprises triphenylphosphine bromide.
- 15 26. The process of claim 1, wherein said silyl protecting group is selectively cleaved off by alkaline hydrolysis and the product of said hydrolysis is sulfonylated.
27. The process of claim 26, wherein the sulfonylation is performing with sulfonylchloride and a base.
- 20 28. The process of claim 1, wherein said silyl protecting group is converted to a thioether using the combination of triphenylphosphine, a dialkyl azodicarboxylate, and a thiol.
29. The process of claim 28, wherein said thiol comprises ethanthiol.
30. The process of claim 28, further comprising the step of reducing said

thioether to leave a methyl group at the C-3' position on the neopterin side chain.

31. The process of claim 30, wherein said reduction is performed with a Raney-Nickel reagent combined with hydrogen.

32. The process of claim 31, wherein said reduction is carried out in an ethanol solvent medium.

33. The process of claim 32, wherein the weight ratio of said ethanol solvent medium to said thioether is about 4:1.

34. The process of claim 1, wherein said step (d) is performed with sodium borohydride, and wherein said reduction is performed in dimethylsulfoxide.

35. The process of claim 1, further comprising the step of forming a salt of tetrahydrobiopterin.


36. The process of claim 35, wherein said salt of tetrahydrobiopterin comprises a dichloride salt of tetrahydrobiopterin.

37. The process of claim 36, further comprising the step of purifying said dichloride salt of tetrahydrobiopterin with recrystallization.

38. The process of claim 1, wherein said steps (a) to (e) are carried out in a polar solvent.

39. The process of claim 38, wherein said polar solvent comprises dimethylformamide.

40. A compound of Formula 6.




6

wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, single linear chain alkyl substituted alkylaminomethyleneimine groups, or single branched chain alkyl substituted alkylaminomethyleneimine groups, double linear chain alkyl substituted alkylaminomethyleneimine groups, double branched chain alkyl substituted alkylaminomethyleneimine groups.

41. The compound of claim 40, wherein R1 is selected from the group consisting of N,N-diethylformamideacetal, and N,N-dimethylformamideacetal.

42. A compound of Formula 7



7


wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, and 2,2-dimethylpropanamide; and wherein R2 is a silyl group that is stable under acidic conditions.

43. The compound of claim 42, wherein R1 comprises N,N-dimethylaminomethylene amino, and R2 is selected from the group consisting of

- diethylisopropylsilyl, dimethylisopropylsilyl, dimethylphenylsilyl,
 diphenylisopropoxysilyl, diphenyl-t-butoxysilyl, di-t-butylmethylsilyl, di-t-
 butylsilylene, methyldiisopropylsilyl, methyldiphenylsilyl, t-butylmethoxyphenylsilyl,
 t-butyltrimethylsilyl, hexyldimethylsilyl, triethylsilyl, 1,1,3,3,-tetra-
 5 isopropylidisiloxane, triisopropylsilyl, trimethylsilyl, trimethylsilyloxycabomyl, and t-
 butyldiphenylsilanoyl.

44. The compound of claim 43, wherein R2 comprises t-
 butyldiphenylsilanoyl.

45. A compound of Formula 7a




wherein R2 is a silyl group that is stable under acidic conditions.

46. The compound of claim 45, wherein R2 is selected from the group
 consisting of diethylisopropylsilyl, dimethylisopropylsilyl, dimethylphenylsilyl,
 diphenylisopropoxysilyl, diphenyl-t-butoxysilyl, di-t-butylmethylsilyl; di-t-
 15 butylsilylene, methyldiisopropylsilyl, methyldiphenylsilyl, t-butylmethoxyphenylsilyl,
 t-butyltrimethylsilyl, hexyldimethylsilyl, triethylsilyl, 1,1,3,3,-tetra-
 isopropylidisiloxane, triisopropylsilyl, trimethylsilyl, trimethylsilyloxycabomyl, and t-
 butyldiphenylsilanoyl.

47. The compound of claim 46, wherein R2 comprises t-
 20 butyldiphenylsilane.

48. A compound of Formula 8



8


wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R2 is a silyl group that is stable under acidic conditions; and R4 a substituted acetal or ketal group that is stable under alkaline conditions.

49. The compound of claim 48, wherein said substituted acetal or ketal group is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals.

50. The compound of claim 49, wherein said substituted acetal or ketal group is selected from the group consisting of methylene acetal, ethylidene acetal, t-butylmethylidene ketal, 1-t-butylethylidene ketal, 1-phenylethylidene ketal, 1-(4-methoxyphenyl)ethylidene acetal, 2,2,2-trichloroethylidene acetal, acrolein acetal, cyclopentylidene ketal, cyclohexylidene ketal, cycloheptylidene ketal, benzylidene acetal, p-methoxybenzylidene acetal, 2,4-dimethoxybenzylidene ketal, 3,4-dimethoxybenzylidene acetal, 2-nitrobenzylidene acetal, 4-nitrobenzylidene acetal, mesitylene acetal, 1-naphthaldehyde acetal, benzophenone ketal, and isopropylideneketal.

51. The compound of claim 48, wherein R3 comprises an N,N-dimethylaminomethylene substituted amino group.

52. A compound of Formula 9.



9


5 wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and wherein R5 is a halogen.

53. The compound of claim 52, wherein R5 is selected from the group consisting of chlorine, bromine, and iodine.

15 54. The compound of claim 52, wherein R4 comprises dimethylacetal.

55. The compound of claim 52, wherein R3 comprises a N,N-dimethylaminomethylene substituted amino group.

56. A compound of Formula 10.




10

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; and R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals.

57. The compound of claim 56, wherein R4 comprises dimethylacetal.

58. The compound of claim 56, wherein R3 comprises a N,N-dimethylaminomethylene substituted amino group.


59. A compound of Formula 11.



11

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; and R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals.

60. A compound of Formula 12.



12

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and R6 is selected from the group consisting of linear chain alkyl substituted sulfonates, branched chain alkyl substituted sulfonates, and aryl substituted sulfonates.


61. The compound of claim 60, wherein R4 comprises dimethylacetal.

62. The compound of claim 60, wherein R3 comprises a N,N-

15 dimethylaminomethylene substituted amino group.

63. The compound of claim 60, wherein R6 comprises a tosyl group.

64. A compound of Formula 11a



11a

wherein R3 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R4 is selected from the group consisting of linear alkyl substituted acetals or ketals, branched alkyl chain substituted acetals or ketals, and aryl substituted acetals or ketals; and R7 is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

10 65. A process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof, comprising the following steps:

(a) reacting pterin at the C-6 position to prepare a 6-substituted pterin;

(b) protecting the primary amine group at C-2 of neopterin with a 2-amino protecting group;

15 (c) metalation of the protected 6-substituted pterin;

(d) coupling of the product of the metalation of the protected 6-substituted pterin with lactic acid or a precursor of lactic acid;

(e) removing the 2-amino protecting group; and

(f) erythro-selective reduction.

20 66. The process of claim 65, wherein the steps (e) and (f) are carried in the same reaction vessel.

67. The process of claim 65, wherein the product from step (e) is isolated.

68. The process of claim 65, wherein the 6-substituted pterin comprises a 6-halogenated pterin.

69. The process of claim 68, wherein said 6-halogenated pterin is selected from the group consisting of 6-chloropterin, 6-bromopterin, and 6-iodopterin.

5 70. The process of claim 69, wherein said 6-halogenated pterin comprises 6-iodopterin.

71. The process of claim 65, wherein the 6-substituted pterin comprises 6-sulfonated pterin.

72. The process of claim 65, wherein said 2-amino protecting group is
10 selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, linear chain alkyl single substituted
15 amido groups, branched chain alkyl single substituted amido groups, and aryl substituted amido groups.

73. The process of claim 72, wherein said 2-amino protecting group comprises a dialkylformamidedialkylacetal.

74. The process of claim 73, wherein said dialkylformamidedialkylacetal is
20 selected from the group consisting of N,N-dimethylformamidediethylacetal, N,N-dimethylformamidedimethylacetal, and bis-dimethylamino-alkoxymethane.

75. The process of claim 72, wherein said 2-amino protecting group is selected from the group consisting of linear chain alkyl single substituted amido groups, branched chain alkyl single substituted amido groups, aryl substituted amido

group, a pivaloyl group, and 2,2-dimethylpropanamido.

76. The process of claim 72, wherein said 2-amino protecting group comprises a pivaloyl group.

77. The process of claim 65, wherein said metalation is performed with a reagent selected from the group consisting of RMgX, alkyl-metal complexes, and metals, wherein X is a halogen, and R is selected from the group consisting of alkyl groups, and aryl groups.

78. The process of claim 77, wherein said metalation is performed with a RMgX reagent, and wherein R is an isopropyl group and X is chloride.

79. The process of claim 77, wherein said alkyl-metal complex comprises an alkyl-metallic lithium complex.

80. The process of claim 79, wherein said alkyl-metallic lithium complex is selected from the group consisting of n-butyllithium, and t-butyllithium.

81. The process of claim 65, wherein said coupling is performed between said protected 6-metalated pterin and a protected lactic acid chloride.

82. The process of claim 81, wherein said protected lactic acid chloride comprises a hydroxyl protected lactic acid chloride.

83. The process of claim 82, wherein said hydroxyl protected lactic acid chloride comprises 2-acetoxypropionic chloride.

84. The process of claim 65, wherein said precursor of lactic acid is selected from the group consisting of 2-oxopropanoyl chlorides, and 2-oxopropanal.

85. The process of claim 84, further comprising the step of reducing the resulting diketones.

86. The process of claim 65, wherein said metalation and said coupling

steps are carried in the same reaction vessel.

87. The process of claim 86, wherein said metalation and said coupling steps are performed with sonification.

88. The process of claim 65, wherein said erythro-selective reduction is performed with sodium borohydride in an alkaline medium.


89. The process of claim 65, wherein said erythro-selective reduction is performed with hydrogen and a catalytic amount of platinum dioxide.

90. The process of claim 65, further comprising the step of forming a salt of tetrahydrobiopterin.

91. The process of claim 65, wherein said salt of tetrahydrobiopterin comprises a dihydrochloride salt of tetrahydrobiopterin.

92. The process of claim 90, further comprising the step of crystallization of said salt of tetrahydrobiopterin.

93. A compound of Formula 2




2

wherein X is selected from the group consisting of chlorine, bromine, iodine, and sulfonates; R₁ is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl

substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, single linear

chain alkyl substituted alkylaminomethylene-imine groups, single branched chain alkyl substituted alkylaminomethylene-imine groups, double linear chain alkyl substituted alkylaminomethylene-imine groups, and double branched chain alkyl substituted alkylaminomethylene-imine groups; and R2 is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups.


94. A compound of Formula 3



wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups; branched chain alkyl substituted sulfur groups, and 2,2-dimethylpropanamide; R2 is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups; and M is selected from the group consisting of boron, silicon, zirconium, titanium, sodium, aluminum, nickel, cobalt, scandium, chromium, ytterbium, lithium, magnesium, zinc, palladium, copper, manganese, cesium, and tin.

95. The compound of claim 94, wherein R1 comprises an N,N-dimethylaminomethylene substituted amino group.


96. A compound of Formula 4



wherein R1 is selected from the group consisting of NH₂, 2,2-dimethylpropanamide, single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, and branched chain alkyl substituted sulfur groups; R2 is selected from the group consisting of hydrogen, linear chain alkyl groups, branched chain alkyl groups, and aryl groups; and R3 is an acyl group.

97. The compound of claim 96, wherein R1 comprises an N,N-dimethylaminomethylene substituted amino group.

98. A compound of Formula 5



wherein R3 is an acyl group.

99. A process for forming enantiomerically-enriched tetrahydrobiopterin or a salt thereof from neopterin, comprising the following steps:

(a) protecting the primary amine group at C-2 of neopterin with a 2-amino protecting group;

(b) converting the primary hydroxyl group of neopterin to a thioether; and

(c) reduction of the thioether leaving a methyl group at the C-3' position on the neopterin side chain.

100. The process of claim 99, wherein said step (c) results in the removal of the 2-amino protecting group, and an erythro-selective reduction to yield
5 tetrahydrobiopterin.

101. The process of claim 100, wherein said step (c) is performed with the use of a reducing agent comprising Raney-Nickel and hydrogen.

102. The process of claim 101, wherein said step (c) is performed at a temperature above 50 degrees Celsius.

103. The process of claim 102, wherein said step (c) is performed in a polar
10 aprotic solvent.

104. The process of claim 103, wherein said polar aprotic solvent comprises ethanol.

105. The process of claim 99, wherein said step (c) does not result in the
15 removal of said 2-amino protecting group and does not result in an erythro-selective reduction.

106. The process of claim 105, further comprising the step of removal of removal of the 2-amino protecting group.

107. The process of claim 106, wherein said removal of the 2-amino
20 protecting group is performed with zinc dichloride in an ethanol solvent medium.

108. The process of claim 105, further comprising the step of an erythro-selective reduction to yield tetrahydrobiopterin.

109. The process of claim 108, wherein said erythro-selective reduction is performed with sodium borohydride in an alkaline medium.

110. The process of claim 109, wherein said erythro-selective reduction is performed with hydrogen and a catalytic amount of platinum dioxide.

111. The process of claim 105, wherein said step (c) is performed with the use of a reducing agent comprising Raney-Nickel.

5 112. The process of claim 111, wherein said step (c) is performed at room temperature.

113. The process of claim 112, wherein said step (c) is performed in a polar aprotic solvent.

114. The process of claim 113, wherein said polar aprotic solvent comprises
10 ethanol.

115. The process of claim 99, further comprising the step of forming a salt of tetrahydrobiopterin.

116. The process of claim 115, wherein said salt of tetrahydrobiopterin comprises a dichloride salt of tetrahydrobiopterin.

15 117. The process of claim 116, further comprising the step of purifying said dichloride salt of tetrahydrobiopterin with recrystallization.

118. The process of claim 99, wherein said 2-amino protecting group is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, single substituted amino groups, double substituted amino groups, aryl single
20 substituted amino groups, linear alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, linear alkyl single substituted alkylaminomethylene-imine groups, linear alkyl double substituted alkylaminomethylene-imine groups, branched alkyl single substituted alkylaminomethylene-imine groups, and branched alkyl double substituted alkylaminomethylene-imine groups.

119. The process of claim 118, wherein said 2-amino protecting group comprises a dialkylformamidedialkylacetal group.

120. The process of claim 119, wherein said dialkylformamidedialkylacetal group is selected from the group consisting of N,N-dimethylformamidediethylacetal, and N,N-dimethylformamidedimethylacetal.

121. The process of claim 118, wherein said 2-amino protecting group comprises a pivaloyl derivative of neopterin.


122. The process of claim 121, wherein said protecting the 2-amino group of neopterin is prepared by a two step process comprising, preparing a tetrapivaloyl derivative of neopterin, and performing alkaline hydrolysis on three pivaloyl groups on said tetrapivaloyl derivative of neopterin.

123. The process of claim 99, wherein said converting the primary hydroxyl group is performed using a disulfide reagent and a trialkylphosphine reagent.

124. The process of claim 123, wherein said disulfide reagent comprises diphenyl disulfide.

125. The process of claim 123, wherein said trialkylphosphine reagent comprises tributylphosphine.

126. A compound of Formula 15



15

wherein R1 is selected from the group consisting of single linear chain alkyl substituted amino groups, single branched chain alkyl substituted amino groups, double linear chain alkyl substituted amino groups, double branched chain alkyl

- substituted amino groups, aryl single substituted amino groups, linear chain alkyl substituted sulfur groups, branched chain alkyl substituted sulfur groups, single linear chain alkyl substituted alkylaminomethyleneimine groups, or single branched chain alkyl substituted alkylaminomethyleneimine groups, double linear chain alkyl substituted alkylaminomethyleneimine groups, double branched chain alkyl substituted alkylaminomethyleneimine groups; and R2 is selected from the group consisting of linear chain alkyl groups, branched chain alkyl groups, and aryl groups.

127. The compound of claim 126, wherein R1 comprises a dialkylalkylaminomethyleneimine group.

128. The compound of claim 127, wherein R1 comprises dimethylaminomethyleneimine.

129. The compound of claim 126, wherein R2 comprises benzene.




Figure 1




Figure 2




Figure 3




Figure 4